

In this problem the disk method was simpler.

**EXAMPLE 4** □ Find the volume of the solid obtained by rotating the region bounded by  $y = x - x^2$  and  $y = 0$  about the line  $x = 2$ .





FIGURE 10

**SOLUTION** Figure 10 shows the region and a cylindrical shell formed by rotation about the line  $x = 2$ . It has radius  $2 - x$ , circumference  $2\pi(2 - x)$ , and height  $x - x^2$ . The volume of the given solid is


$$\begin{aligned} V &= \int_0^1 2\pi(2 - x)(x - x^2) dx = 2\pi \int_0^1 (x^3 - 3x^2 + 2x) dx \\ &= 2\pi \left[ \frac{x^4}{4} - x^3 + x^2 \right]_0^1 = \frac{\pi}{2} \end{aligned}$$

### 6.3 Exercises

1. Let  $S$  be the solid obtained by rotating the region shown in the figure about the  $y$ -axis. Explain why it is awkward to use slicing to find the volume  $V$  of  $S$ . Sketch a typical approximating shell. What are its circumference and height? Use shells to find  $V$ .



2. Let  $S$  be the solid obtained by rotating the region shown in the figure about the  $y$ -axis. Sketch a typical cylindrical shell and find its circumference and height. Use shells to find the



volume of  $S$ . Do you think this method is preferable to slicing? Explain.

- 3-7 □ Use the method of cylindrical shells to find the volume generated by rotating the region bounded by the given curves about the  $y$ -axis. Sketch the region and a typical shell.

3.  $y = 1/x$ ,  $y = 0$ ,  $x = 1$ ,  $x = 2$
4.  $y = x^2$ ,  $y = 0$ ,  $x = 1$
5.  $y = e^{-x^2}$ ,  $y = 0$ ,  $x = 0$ ,  $x = 1$
6.  $y = x^2 - 6x + 10$ ,  $y = -x^2 + 6x - 6$
7.  $y^2 = x$ ,  $x = 2y$

8. Let  $V$  be the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = \sqrt{x}$  and  $y = x^2$ . Find  $V$  both by slicing and by cylindrical shells. In both cases draw a diagram to explain your method.

- 9-14 □ Use the method of cylindrical shells to find the volume of the solid obtained by rotating the region bounded by the given curves about the  $x$ -axis. Sketch the region and a typical shell.

9.  $x = 1 + y^2$ ,  $x = 0$ ,  $y = 1$ ,  $y = 2$

10.  $x = \sqrt{y}$ ,  $x = 0$ ,  $y = 1$

- 11.  $y = x^2, y = 9$
- 12.  $y^2 - 6y + x = 0, x = 0$
- 13.  $y = \sqrt{x}, y = 0, x + y = 2$
- 14.  $x + y = 3, x = 4 - (y - 1)^2$


15–20 □ Use the method of cylindrical shells to find the volume generated by rotating the region bounded by the given curves about the specified axis. Sketch the region and a typical shell.

- 15.  $y = x^2, y = 0, x = 1, x = 2$ ; about  $x = 1$
- 16.  $y = x^2, y = 0, x = -2, x = -1$ ; about the  $y$ -axis
- 17.  $y = x^2, y = 0, x = 1, x = 2$ ; about  $x = 4$
- 18.  $y = 4x - x^2, y = 8x - 2x^2$ ; about  $x = -2$
- 19.  $y = \sqrt{x - 1}, y = 0, x = 5$ ; about  $y = 3$
- 20.  $y = x^2, x = y^2$ ; about  $y = -1$

21–26 □ Set up, but do not evaluate, an integral for the volume of the solid obtained by rotating the region bounded by the given curves about the specified axis.

- 21.  $y = \ln x, y = 0, x = 2$ ; about the  $y$ -axis
- 22.  $y = x, y = 4x - x^2$ ; about  $x = 7$
- 23.  $y = x^4, y = \sin(\pi x/2)$ ; about  $x = -1$
- 24.  $y = 1/(1 + x^2), y = 0, x = 0, x = 2$ ; about  $x = 2$
- 25.  $x = \sqrt{\sin y}, 0 \leq y \leq \pi, x = 0$ ; about  $y = 4$
- 26.  $x^2 - y^2 = 7, x = 4$ ; about  $y = 5$

- 27. Use the Midpoint Rule with  $n = 4$  to estimate the volume obtained by rotating about the  $y$ -axis the region under the curve  $y = \tan x, 0 \leq x \leq \pi/4$ .
- 28. If the region shown in the figure is rotated about the  $y$ -axis to form a solid, use the Midpoint Rule with  $n = 5$  to estimate the volume of the solid.



29–32 □ Each integral represents the volume of a solid. Describe the solid.

- 29.  $\int_0^{\pi/2} 2\pi x \cos x \, dx$
- 30.  $\int_0^9 2\pi y^{3/2} \, dy$
- 31.  $\int_0^1 2\pi(x^3 - x^7) \, dx$
- 32.  $\int_0^\pi 2\pi(4 - x) \sin^4 x \, dx$

33–34 □ Use a graph to estimate the  $x$ -coordinates of the points of intersection of the given curves. Then use this information to estimate the volume of the solid obtained by rotating about the  $y$ -axis the region enclosed by these curves.

- 33.  $y = 0, y = x + x^2 - x^4$
- 34.  $y = x^4, y = 3x - x^3$

35–40 □ The region bounded by the given curves is rotated about the specified axis. Find the volume of the resulting solid by any method.

- 35.  $y = x^2 + x - 2, y = 0$ ; about the  $x$ -axis
- 36.  $y = x^2 - 3x + 2, y = 0$ ; about the  $y$ -axis
- 37.  $y = 5, y = x + (4/x)$ ; about  $x = -1$
- 38.  $x = 1 - y^4, x = 0$ ; about  $x = 2$
- 39.  $x^2 + (y - 1)^2 = 1$ ; about the  $y$ -axis
- 40.  $x^2 + (y - 1)^2 = 1$ ; about the  $x$ -axis

41–43 □ Use cylindrical shells to find the volume of the solid.

- 41. A sphere of radius  $r$
- 42. The solid torus of Exercise 59 in Section 6.2
- 43. A right circular cone with height  $h$  and base radius  $r$
- 44. Suppose you make napkin rings by drilling holes with different diameters through two wooden balls (which also have different diameters). You discover that both napkin rings have the same height  $h$ , as shown in the figure.
  - (a) Guess which ring has more wood in it.
  - (b) Check your guess: Use cylindrical shells to compute the volume of a napkin ring created by drilling a hole with radius  $r$  through the center of a sphere of radius  $R$  and express the answer in terms of  $h$ .

